


# Association of germline variation with the survival of women with *BRCA1/2* pathogenic variants and breast cancer


Taru A. Murañen, Sofia Khan, Rainer Fagerholm, Kristiina Aittomäki, Julie M. Cunningham, Joe Dennis, Goska Leslie, Lesley McGuffog, Michael T. Parsons, Jacques Simard, Susan Slager, Penny Soucy, Douglas F. Easton, Marc Tischkowitz, Amanda B. Spurdle, kConFab Investigators, Rita K. Schmutzler, Barbara Wappenschmidt, Eric Hahnen, Maartje J. Hooning, HEBON Investigators, Christian F. Singer, Gabriel Wagner, Mads Thomassen, Inge Sokilde Pedersen, Susan M. Domchek, Katherine L. Nathanson, Conxi Lazaro, Maria Rossing, Irene L. Andrulis, Manuel R. Teixeira, Paul James, Judy Garber, Jeffrey N. Weitzel, SWE-BRCA Investigators, Anna Jakubowska, Drakoulis Yannoukakos, Esther M. John, Melissa C. Southey, Marjanka K. Schmidt, Antonis C. Antoniou, Georgia Chenevix-Trench, Carl Blomqvist, Heli Nevanlinna

## Supplementary Information

Supplementary Figure 1. The survival effect associated with rs57025206 in all <i>BRCA1</i> carriers.....	2
Supplementary Figure 2. Manhattan plots.....	3
Supplementary Figure 3. Plots for survival variants discovered in <i>BRCA1</i> carriers.....	4
Supplementary Figure 4. Plots for survival variants discovered in the meta-analysis.....	5
Supplementary Figure 5. Plots for survival variants discovered in <i>BRCA2</i> carriers.....	6
Supplementary Table 1. Description of CIMBA studies.....	7
Supplementary Table 2. Additional models for <i>BRCA1</i> carriers.....	8
Supplementary Table 3. Additional meta-analysis models.....	8
Supplementary Table 4. Additional models for <i>BRCA2</i> carriers.....	9
Supplementary Table 5. Survival associations of variants with age-dependent survival effect in <i>BRCA2</i> carriers, stratified by the tumor ER-status.....	9
Supplementary Table 6. Literature-based functional annotation of the target genes.....	10
Supplementary Table 7. Survival associations in the Breast Cancer Association Consortium (BCAC) data.....	14
Supplementary References.....	15
Supplementary note.....	19
Supplementary Data 1. Identification of potential target genes based on variant position and functional annotations.....	21
Supplementary Data 2. eQTL-analysis in GTEx and Westra et al. data.....	21




**Supplementary Figure 1.** The survival effect associated with rs57025206 in all *BRCA1* carriers.  
a) Kaplan-Meier plot stratified by rs57025206. b) Forest plot of hazard ratios (HR) accross country groups.  
P.het: P-value against between-study heterogeneity.



**Supplementary Figure 2. Manhattan plots.**


Manhattan plots from survival-association analyses of a) *BRCA1* carriers, b) *BRCA1* carriers with ER-negative breast cancer, c) *BRCA2* carriers, d) *BRCA2* carriers from variant-diagnosis age interaction analysis, and e) a meta-analysis of nominal survival associations in *BRCA1* and *BRCA2* carriers. P-values from likelihood ratio test plotted against genomic position. (Green dashed line:  $5 \cdot 10^{-8}$ ; blue dashed line:  $5 \cdot 10^{-7}$ )



**Supplementary Figure 3.** Plots for survival variants discovered in *BRCA1* carriers


Kaplan-Meier curves graphically presenting the proportion of surviving patients during the 15 years after primary breast cancer diagnosis in the discovery data (see Table 2), stratified by the 10 *BRCA1*-associated survival variants (1st and 3rd panel from left). Forest plots presenting the effect sizes in different country groups (2nd and 4th panel). Because the carriers of effect alleles of rs1829314 and rs78150318 had no events, the forest plots could not be drawn.





**Supplementary Figure 4.** Plots for survival variants discovered in the meta-analysis

Kaplan-Meier curves stratified by the four variants with consistent survival effects for *BRCA1* and *BRCA2* carriers (1st and 3rd panel from the left, respectively). Forest plots presenting the effect sizes in different country groups (2nd and 4th panel).



**Supplementary Figure 5.** Plots for survival variants discovered in *BRCA2* carriers

Kaplan-Meier curves stratified by the four variants with age-dependent survival effect for *BRCA2* carriers in three age-groups: patients diagnosed before the age of 40 years (left), between the age of 40 and age of 50 years (middle), and after the age of 50 years (right).

**Supplementary Table 1.** Description of CIMBA studies.

Study acronym	Study name	Country	Country group	Study subject ascertainment	Number of study subjects		Number of events	
					BRCA1	BRCA2	BRCA1	BRCA2
BCFR-AU	Australian site of the Breast Cancer Family Registry	Australia	Australia	Population based	25		8	
BCFR-NC	Northern California site of the Breast Cancer Family Registry	USA	USA	Population based	31		6	
BCFR-ON/ OCGN	Ontario site of the Breast Cancer Family Registry/ Ontario Cancer Genetics Network	Canada	Canada	Clinic and population based	73	53	15	17
CBCS	Copenhagen Breast Cancer Study	Denmark	Denmark	Clinic based	73	63	14	9
COH	City of Hope Cancer Center	USA	USA	Clinic based	101		15	
DEMOKRITOS	National Centre for Scientific Research Demokritos	Greece	Balkan	Clinic based	57		9	
DFCI	Dana Farber Cancer Institute	USA	USA	Clinic based	64	46	8	8
EMBRACE	Epidemiological Study of Familial Breast Cancer	UK	UK	Clinic based	694	678	95	89
GC-HBOC	German Familial Breast Group	Germany	Central-Europe	Clinic based	368	207	33	17
HEBCS	Helsinki Breast Cancer Study	Finland	Scandinavia	Clinic based	62	61	18	16
HEBON	Hereditary Breast and Ovarian cancer study the Netherlands	Netherlands	Netherlands	Clinic based	234	93	41	28
ICO	Institut Català d'Oncologia	Spain	Iberia	Clinic based	87	100	14	16
IHCC	International Hereditary Cancer Centre	Poland	East-Europe	Clinic based	94		7	
IPOBCS	Portuguese Oncology Institute-Porto Breast Cancer Study	Portugal	Iberia	Clinic based	39	82	12	11
KCONFAB	Kathleen Cuningham Consortium for Research into Familial Breast Cancer	Australia	Australia	Clinic based	331	245	53	37
MUV	General Hospital Vienna	Austria	Central-Europe	Clinic based	218	108	38	17
OUH	Odense University Hospital	Denmark	Denmark	Clinic based	172	143	30	27
SWE-BRCA	Swedish Breast Cancer Study	Sweden	Scandinavia	Clinic based	97		19	
UPENN	University of Pennsylvania	USA	USA	Clinic based	118	82	13	11
VFCTG	Victorian Familial Cancer Trials Group	Australia	Australia	Clinic based	70	48	13	8



**Supplementary Table 2.** Additional models for BRCA1 carriers.

Variant effects in multivariate and in breast cancer-specific survival analyses for the SNPs, which were considered novel discoveries in the analysis of *BRCA1* carriers and *BRCA1* carriers with ER-negative breast cancer.

SNP	Analysis subgroup	Genetic model	HR for all-cause mortality in a model adjusted for tumor characteristics		Nominal HR for breast cancer-specific death		HR for breast cancer-specific death in a model adjusted for tumor characteristics	
			HR	[95% CI]	HR	[95% CI]	HR	[95% CI]
rs12126206	BRCA1 all BC	dominant	1.96	[1.37-2.81]	1.53	[1.08-2.17]	1.91	[1.07-3.39]
rs5028286	BRCA1 all BC	dominant	1.55	[1.11-2.17]	1.44	[1.04-1.99]	1.21	[0.69-2.10]
rs736418	BRCA1 all BC	dominant	0.60	[0.43-0.83]	0.72	[0.53-0.96]	0.41	[0.24-0.70]
rs147857072	BRCA1 all BC	dominant	4.95	[2.67-9.18]	2.55	[1.41-4.61]	3.32	[1.28-8.58]
rs59010985	BRCA1 all BC	per-allele linear	0.49	[0.34-0.70]	0.61	[0.44-0.83]	0.66	[0.40-1.09]
rs34987256†	BRCA1 all BC	per-allele linear	1.44	[1.09-1.89]	1.68	[1.31-2.16]	1.61	[1.04-2.48]
rs1829314	BRCA1 ER-negative	dominant	0.00	[0.00-0.00]	0.00	[0.00-0.00]	0.00	[0.00-0.00]
rs57025206	BRCA1 ER-negative	per-allele linear	6.19	[3.73-10.3]	4.68	[2.81-7.80]	4.98	[1.91-13.0]
rs11245414	BRCA1 ER-negative	per-allele linear	0.68	[0.51-0.91]	0.54	[0.40-0.72]	0.63	[0.41-0.97]
rs78150318	BRCA1 ER-negative	dominant	0.00	[0.00-0.00]	0.00	[0.00-0.00]	0.00	[0.00-0.00]

†The hazard associated with the SNP did not significantly violate of the proportional hazards assumption in the multivariate models or in the analyses of breast cancer-associated death.

**Supplementary Table 3.** Additional meta-analysis models.

Variant effects in multivariate and in breast cancer-specific survival analyses for the SNPs, which were considered novel discoveries in the *BRCA1-BRCA2*-carrier meta-analysis.

SNP	Analysis subgroup	Genetic model	HR for all-cause mortality in a model adjusted for tumor characteristics		Nominal HR for breast cancer-specific death		HR for breast cancer-specific death in a model adjusted for tumor characteristics	
			HR	[95% CI]	HR	[95% CI]	HR	[95% CI]
rs551383190	BRCA1/BRCA2	per-allele linear	0.61	[0.46-0.82]	0.63	[0.49-0.81]	0.68	[0.50-0.94]
rs117422049	BRCA1/BRCA2	per-allele linear	3.43	[0.83-14.3]	3.24	[1.97-5.32]	3.60	[2.00-6.47]
rs4879914	BRCA1/BRCA2	per-allele linear	1.40	[1.18-1.67]	1.39	[1.20-1.60]	1.49	[1.24-1.80]
rs2320070	BRCA1/BRCA2	per-allele linear	1.31	[1.06-1.63]	1.40	[1.17-1.68]	1.46	[1.02-2.09]



**Supplementary Table 4.** Additional models for BRCA2 carriers.

Variant effects in multivariate and in breast cancer-specific survival analyses for the SNPs, which had age-dependent survival effect for *BRCA2* carriers.

SNP	Analysis subgroup	Genetic model	HR for all-cause mortality in a model adjusted for tumor characteristics				Nominal HR for breast cancer-specific death				HR for breast cancer-specific death in a model adjusted for tumor characteristics			
			dgAge < 40 years		dgAge ≥ 40		dgAge < 40 years		dgAge ≥ 40		dgAge < 40 years		dgAge ≥ 40	
			HR	[95% CI]	HR	[95% CI]	HR	[95% CI]	HR	[95% CI]	HR	[95% CI]	HR	[95% CI]
rs372812554	BRCA2 all BC	per-allele interaction with age	2.11	[1.33-3.34]	0.84	[0.57-1.23]	1.91	[1.32-2.76]	0.74	[0.53-1.04]	2.04	[1.09-3.85]	0.66	[0.39-1.12]
rs2109815	BRCA2 all BC	per-allele interaction with age	1.70	[1.09-2.67]	0.92	[0.64-1.32]	1.78	[1.24-2.56]	0.87	[0.64-1.18]	2.01	[1.17-3.46]	1.12	[0.66-1.91]
rs35431863	BRCA2 all BC	per-allele interaction with age	0.58	[0.29-1.14]	1.29	[0.94-1.78]	0.55	[0.38-0.78]	1.27	[0.98-1.65]	0.62	[0.27-1.44]	1.40	[0.94-2.09]
rs11255420	BRCA2 all BC	per-allele interaction with age	0.62	[0.32-1.18]	1.28	[0.87-1.90]	0.67	[0.44-1.01]	1.73	[1.28-2.33]	0.44	[0.19-1.05]	1.29	[0.74-2.24]

**Supplementary Table 5.** Survival associations of variants with age-dependent survival effect in BRCA2 carriers, stratified by the tumor ER-status.

SNP	ER-positive BRCA2 carriers (n:1067)		ER-negative BRCA2 carriers (n: 302)	
	under 40 years HR [95% CI]	over 40 years HR [95% CI]	under 45 years HR [95% CI]	over 45 years HR [95% CI]
rs372812554	1.67 [1.19 - 2.36]	0.69 [0.49 - 0.98]	1.43 [0.50 - 4.05]	1.40 [0.77 - 2.54]
rs2109815	1.86 [1.24 - 2.79]	0.75 [0.56 - 1.01]	1.57 [0.84 - 2.93]	0.46 [0.20 - 1.05]
rs35431863	0.54 [0.37 - 0.80]	1.48 [1.15 - 1.90]	0.97 [0.47 - 2.02]	1.31 [0.71 - 2.41]
rs11255420	0.59 [0.41 - 0.83]	1.55 [1.24 - 1.93]	0.47 [0.15 - 1.46]	0.86 [0.43 - 1.69]

**Supplementary Table 6.** Literature-based functional annotation of the target genes.

Gene	Function		Ref.	
KIF26B	silencing in breast cancer cell lines	reduces colony-formation	1, 2	
		reduces migration and invasion		
		leads to increased E-cadherin expression and reduced N-cadherin expression		
		induces apoptosis		
		aggregates cells in G0/G1 phase		
	overexpression in breast cancer cell lines	increases proliferation (increased cell count)	2	
		promotes migration and invasion		
		positively affects FGF2 expression and secretion		
		leads to MAPK/ERK pathway activation		
	knock-down in a xenograft model	reduces tumor formation	1, 2	
		reduces the frequency of lung metastases		
		mRNA expression in mammary tumors		
		protein expression in mammary tumors		2
mRNA expression in breast cancer cell lines		1		
high expression in mammary tumors		3		
enhances microtubule stabilization		4		
SGCZ	loss in mammary tumors	is more frequent in early-onset tumors (METABRIC)	6, 7	
		is associated with poor prognosis		
		does not affect SGCZ mRNA expression in mammary tumors, which is generally very low		
	MIR383	is located in intron 1-2 of SGCZ	8	
		overexpression in mouse ovarian cells		
	overexpression in ovarian cancer cell lines	enhances estradiol release from ovaries via repression of RMBS1, and consecutively also MYC	9	
		sensitizes the cells to paclitaxel		
	RALGDS	is recruited to endosomal compartments	by RILP	10
		mediates the RILP-dependent	inhibition of cell proliferation, migration and invasion	
		serves	as a guanine nucleotide exchange factor for Ral	
interaction with RILP		affects the MAPK/ERK pathway		
does not directly stimulate		MAPK/ERK	11	
induces cytoskeletal reorganization		in response to FPR1 stimulation	12	
MBIP		was the top-ranking breast cancer risk gene	in a genome-wide pathway analysis	13
GAS7	is hypomethylated (activated)	in ER-negative breast cancer in comparison to ER-positive breast cancer	14, 15	
	transcript variant B mRNA expression	is low in mammary tumors in comparison to adjacent normal	16	
		is low in early-onset tumors in comparison to late-onset tumors		
		is low in many breast cancer cell lines		
	overexpression in several breast cancer cell lines	reduces proliferation, migration and invasion		
	silencing in MCF-7 breast cancer cell line	increases proliferation, migration and invasion		
	high expression	inhibits actin polymerization via GAS7 – CYFIP1 – Rac1 – WAVE2 complex		
	is regulated by TP53			
	transcript variant B regulatory region	co-localizes with the survival variants (Supplementary Table 5)		
	rs59010985 allele T	is associated with high GAS7 expression (Table 5, Supplementary Table 6)		
	high mRNA expression is associated with	good prognosis of ER-negative breast cancer patients HR = 0.46 [0.37 - 0.58], P = 2.5E-12 (FDR 1%)	17	
		good prognosis of breast cancer patients irrespective of ER-status	7	
		HR per unit fold change in all 0.81 [0.72 – 0.90], P = 1.4E-4, in METABRIC data		
mediates the apoptotic effects of	platinum compounds in hepatocellular carcinoma and neuroblastoma cells	18, 19		
	gefitinib in non-small cell lung carcinoma	20		

Gene	Function		Ref.
CHST9	variant rs1436904	is significantly associated with breast cancer risk in an international consortium study	21
		is associated with survival of triple-negative breast cancer patients in Chinese population	22
		is linked with the survival variant rs537497819 with D'=0.1939, R2=0.0188	
DCAF1	is an E3 ubiquitin ligase substrate receptor	which brings together the substrate and E2 ubiquitin conjugating ligase	23
	recognizes the substrate for ubiquitination	leading to proteasomal degradation of the substrate	
	substrates include	TP53 tumor suppressor, apoptosis regulator	
		ER-alpha via LATS1 interaction	24
		non-histone proteins monomethylated by EZH2 (component of the polycomb repressor complex)	25
	silencing in MCF-7 breast cancer cell line	reduces colony formation	
	regulates	DNA replication	23
		cell cycle progression	
		entry to mitosis	
		cell division	
		ER-alpha level	24
		T-cell activation induced proliferation	26
		T-cell receptor gene reorganization	
		centrosome organization by ubiquitylation as a part of the EDVP complex	27
MIR135A1	deletion in mammary tumors is associated with	higher pathologic stage	28
		younger age of onset	
		ductal histologic type	
		worse survival outcome	
		lower expression of pri-miR-135-a-1, primary precursor of mir-135A1	
	expression in mammary tumors	is higher in ER+ than ER- cancers	
	depletion in breast cancer cell lines	increases cell viability, colony formation, migration, and invasion	
		decreases sensitivity to tamoxifen treatment and contributes to acquisition of resistance	
	forced expression in breast cancer cell lines	enhances the activity of MAPK/ERK and PI3K/AKT pathways	
		decreases cell viability, colony formation, migration, and invasion	
		enhances epithelial cellular phenotype	
	level is correlated	suppresses the activity of MAPK/ERK and PI3K/AKT pathways	
		with epithelial markers	
		inversely with mesenchymal markers	
	forced expression in a mouse model	inhibits tumor formation and lung metastasis	
	promoter is bound by ER-alpha		
	directly binds and regulates ER-alpha	creating a negative feedback loop	
	expression promotes resistance to	oxaliplatin in gastric cancer	29
		gefitinib in non-small cell lung cancer	30
3p21.2 (DCAF1 and MIR135A1)	loss in mammary tumors	is more frequent than loss of any other region in 3p	31, 32
		is associated with high grade and ER-, PR- tumors	
ZRANB1	regulates	EZH2 (component of the polycomb repressor complex) by deubiquitination	33
	silencing in triple-negative breast cancer cell lines	reduces proliferation and migration	
		which can be rescued by overexpressing EZH2	
	silencing in a mouse model	suppresses tumor formation and lung metastases	
	silencing in cell lines	increases stress fibers and inhibits migration	34
	expression in normal mammary tissue	is very low	33
	high expression in mammary tumors	is associated with poor survival	
	is required for	inflammatory T-cell response	35

Gene	Function	Ref.
CTBP2	is a transcriptional co-repressor	36
	primes target genes	37
	regulates	36, 38, 39
		p16, CDH1, PTEN
		stem-cell like characteristics of ovarian cancer cells
	silencing in breast cancer cell lines	36, 38, 40
		leads to TP53-dependent cell cycle arrest and apoptosis
		causes cell cycle stasis in aberrant mitosis
		increases the number of binucleate cells and lagging chromosomes
		suppresses cell migration
	overexpression in breast cancer cell lines	38
		increases cell proliferation and cell accumulation in S-phase
		decreases E-cadherin expression
		increases vimentin and MMP2 expression
	overexpression in a mouse xenograft model	
	is expressed ubiquitously in	40
	high expression in mammary tumors	17, 38
	low expression in ovarian tumors	39
RAD51B		and poor prognosis of ovarian cancer patients
	proteasomal degradation can be induced by	40
	may mediate	41
	binds to BRCA1 promoter in ovarian cancer cell lines,	42
	overexpression reduces sensitivity to cisplatin	43, 44
RAD51B	variant rs2588809	21, 45
		is associated with breast cancer risk
		is linked with the survival variant rs78150318 with R2=0.0016, D'=0.4677
RAD51B	silencing is associated with	46, 47
	somatic alterations in BRCA1-defective breast cancer	48
		are associated with poor overall survival of patients
CREB5	is repressed by miR-29c	49
	high expression in ER-positive mammary tumors	17
ASPH	locus is amplified in a subgroup of primary tumors	50
	becomes upregulated in response to	51
	expression is inversely correlated with	
	high expression predicts poor outcome	
	proteins and peptides are highly immunogenic	52, 53
GATA3		IGF1 exposure via MAPK and PI3K/Akt pathways
		tamoxifen sensitivity of breast cancer cell lines
		from endocrine therapy for patients with luminal B breast cancer
		in induced dendritic cells of hepatocellular carcinoma patients
		and may thus be a target for immunotherapy in hepatocellular carcinoma
	is a transcription factor	54-56
	regulates differentiation of	54, 55, 57
	silencing in mammary luminal cells	
		blocks differentiation
		increased proliferation
		leads to abnormalities nuclear size and orientation
		causes cell detachment and increased cell death
	knock-down in mice causes severe defects	54, 55
	modulates ESR1 binding profile	56
	high expression in mammary tumors	58-62
		is associated with hormone-positive/luminal tumor phenotype and low grade
		is associated with good patient prognosis
		does not affect prognosis
		is associated with shorter locoregional relapse time in premenopausal patients
		is associated with decreased rate of pathologic complete response to neoadjuvant chemotherapy
		is associated with increased probability of GATA3 somatic mutation
	regulates spindle orientation	67
	has diverse roles	68



Gene	Function	Ref.
ZNF644	is a component of G9a/GLP complex	69, 70
	which represses transcription via H3K9 methylation	
	which interacts with polycomb repressor complex	
	which is a part of the replisome, required to prevent replication-associated DNA damage	
	recognizes the DNA target sequence	69
	silencing in vitro	71
	decreased proliferation	
CLASP1	sensitized to replication stress	
	increased DNA damage in replicating cells	
	high expression in mammary tumors	17
	is associated with poor prognosis with HR = 1.43 [1.21 - 1.69], P = 2.2E-5 (FDR 2%)	
	regulates microtubule dynamics during mitosis	72, 73
	localizes to the outer region of kinetochore (korona)	
	near the kinetochore-attached microtubule plus-ends	
NIFK	knock-down in human and murine cells	73
	leads to chromosomal instability	
	is required for invasion through 3D matrix	74
	enhancing compression- resistance of growing microtubules	
	high expression in mammary tumors	17
	is associated with good prognosis with HR = 0.71 [0.63 - 0.80], P = 4.9E-9 (FDR 1%)	
	nucleolar protein interacting with the FHA domain of MKI67	75
NIFK-AS1	overexpression in ling cancer cell lines	
	enhances proliferation, migration, and invasion	
	regulates TCF4/ $\beta$ -catenin	
	via repression of RUNX1 and CK1 $\alpha$	
	high expression in mammary tumors	
	is associated with poor patient survival	
	is phosphorylated by CDK1 and GSK3	76
TFCP2L1	transcription is induced by c-Myc and estrogen	
	knock-down in osteosarcoma cell line	
	induced cell cycle arrest in G1 as a result of ribosomal stress	
	is required for ribosomal RNA maturation	
	especially in ITS1 (internal transcribed spacer 1) processing	
	suppresses macrophage M2 polarization;	77
	overexpression in macrophages	
ARHGEF39	suppressed the proliferation of estrogen-stimulated endometrial cancer cells	
	high expression in mammary tumors	17
	is associated with good prognosis with HR = 0.64 [0.54 - 0.74], P = 1.5E-8 (FDR 1%)	
	is required for pluripotency	78
	of embryonic stem cells	
	together with Zf5, Ctcf, E2f1, and Myc	79
	predicts the targets of polycomb repressor complex	
TPM2	low expression in mammary tumors	17
	is associated with poor prognosis with HR = 1.41 [1.20 - 1.67], P = 3.1E-5 (FDR 5%)	
	overexpression in gastric and non-small cell lung	80, 81
	cancer cell lines induces proliferation, migration, and invasion	
	increases Akt phosphorylation activating the Akt/PI3K pathway	
	increases P38 and ATF2 phosphorylation and activates MAPK pathway	
	increases the cellular levels of Cyclin A2, Cyclin D1, and MMP2	
TPM2	increases Rac1 activation	
	knockdown in gastric and non-small cell lung	
	cancer cell lines represses proliferation, migration, and invasion	
	binds actin	82
	to stabilize microfilaments	
	in involved in cytokinesis, cellular vesicle transport, proliferation, migration, and apoptosis	
	expression is reduced	
TPM2	in breast cancer in comparison to adjacent normal tissue	
	in hypoxic conditions	
	silencing in breast cancer cell lines	
	increases invasion and migration	
	contributes to paclitaxel resistance	
	silencing in HeLa and U2OS cells	83
	induces lysosomal destabilization and lysosomal cell death	
TPM2	sensitizes to cisplatin, siramesine, and etoposide	
	low protein expression in mammary tumors	82
	is associated with poor patient survival	
TPM2	low mRNA expression in mammary tumors	17
	is associated with good survival with HR = 0.64 [0.57 - 0.72], P = 2.9E-14 (FDR 1%)	

Gene	Function	Ref.
GBA2	Bile acid b-glucosidase	breaks down glucocylceramide (outside lysosomes)
	Multi-drug resistant MCF-7	has high levels of glucocylceramide due to high glucocylceramide synthase activity
	Glucocylceramide synthase repression	increases drug sensitivity in multi-drug resistant MCF-7
	Overexpression of GBA2	DOES NOT have the same effect in multi-drug resistant MCF-7
	low expression in mammary tumors	is associated with poor survival with HR = 1.58 [1.35 - 1.85], P = 5.6E-9 (FDR 1%)
RUSC2	silencing leads to accumulation of glucosylceramide	in plasma membrane, affecting actin and microtubule dynamics
	silencing in lung cancer cell lines	reduced directional migration (chemotaxis)
		did not affect random migration (chemokinesis)
		causes defective Golgi orientation for chemotaxis
CD72	is required for EGFR-induced	GIT2 (G protein-coupled receptor kinase interacting ArfGAP 2) phosphorylation
		directional cell migration
	interaction with CD5 is required for	regulatory B cell and regulatory T cell reciprocal stimulation
TRAV	is a membrane-bound receptor for CD100	in T cells
	interaction with CD100	is required for T cell activation

**Supplementary Table 7.** Survival associations in the Breast Cancer Association Consortium (BCAC) data.

Variant	BCAC: analysis group	HR [95% CI]	CIMBA: analysis group	HR [95% CI]	CIMBA: analysis group 2	HR [95% CI]
rs12126206	ER-	0.95 [0.83-1.08]	BRCA1 all BC	1.86 [1.50 - 2.32]		
rs5028286	ER-	1.09 [0.99-1.21]	BRCA1 all BC	1.74 [1.42 - 2.12]		
rs736418	ER-	1.01 [0.93-1.09]	BRCA1 all BC	0.60 [0.49 - 0.73]		
rs147857072	ER-	1.00 [0.67-1.49]	BRCA1 all BC	3.41 [2.32 - 5.00]		
rs59010985	ER-	0.96 [0.87-1.06]	BRCA1 all BC	0.58 [0.47 - 0.72]		
rs537497819			BRCA1 all BC	1.49 [1.27 - 1.74]		
rs1829314	ER-	1.10 [0.92-1.31]	BRCA1 ER-	0 [0 - 0]		
rs57025206	ER-	1.06 [0.86-1.29]	BRCA1 ER-	4.37 [3.03 - 6.30]		
rs11245414	ER-	1.00 [0.93-1.08]	BRCA1 ER-	0.56 [0.46 - 0.70]		
rs78150318	ER-	0.94 [0.78-1.15]	BRCA1 ER-	0 [0 - 0]		
rs551383190			BRCA1/BRCA2	0.61 [0.51 - 0.73]		
rs117422049	All	1.15 [0.98-1.35]	BRCA1/BRCA2	2.80 [1.89 - 4.13]		
rs4879914	All	0.99 [0.96-1.02]	BRCA1/BRCA2	1.30 [1.17 - 1.44]		
rs2320070	All	0.99 [0.95-1.03]	BRCA1/BRCA2	1.39 [1.22 - 1.58]		
rs372812554			BRCA2 under 40y	1.75 [1.33 - 2.31]	BRCA2 over 40y	0.75 [0.59 - 0.96]
rs2109815	ER+	1.03 [0.98-1.08]	BRCA2 under 40y	1.65 [1.24 - 2.19]	BRCA2 over 40y	0.75 [0.59 - 0.95]
rs35431863	ER+	1.01 [0.97-1.06]	BRCA2 under 40y	0.55 [0.43 - 0.71]	BRCA2 over 40y	1.32 [1.09 - 1.60]
rs11255420	ER+	0.98 [0.93-1.03]	BRCA2 under 40y	0.59 [0.41 - 0.83]	BRCA2 over 40y	1.55 [1.24 - 1.93]

## Supplementary References

1. Gu S, Liang H, Qi D, Mao L, Mao G, Qian L, Zhang S. Knockdown of KIF26B inhibits breast cancer cell proliferation, migration, and invasion. *Onco Targets Ther* 11:3195-3203, 2018.
2. Teng Y, Guo B, Mu X, Liu S. KIF26B promotes cell proliferation and migration through the FGF2/ERK signaling pathway in breast cancer. *Biomed Pharmacother* 108:766-773, 2018.
3. Wang Q, Zhao ZB, Wang G, Hui Z, Wang MH, Pan JF, Zheng H. High expression of KIF26B in breast cancer associates with poor prognosis. *PLoS One* 8:e61640, 2013.
4. Guillabert-Gourgues A, Jaspard-Vinassa B, Bats M, Sewduth RN, Franzl N, Peghaire C, Jeanningros S, Moreau C, Roux E, Larrieu-Lahargue F, Dufourcq P, Couffignal T, Dupl  a C. Kif26b controls endothelial cell polarity through the dishevelled/Daam1-dependent planar cell polarity-signaling pathway. *Mol Biol Cell* 27:941-953, 2016.
5. Pu Y, Yi Q, Zhao F, Wang H, Cai W, Cai S. MiR-20a-5p represses multi-drug resistance in osteosarcoma by targeting the KIF26B gene. *Cancer Cell Int* 16:64, 2016.
6. Chi C, Murphy LC, Hu P. Recurrent copy number alterations in young women with breast cancer. *Oncotarget* 9:11541-11558, 2018.
7. Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, Speed D, Lynch AG, Samarajiwa S, Yuan Y, Graf S, Ha G, Haffari G, Bashashati A, Russell R, McKinney S, METABRIC Group, Langerod A, Green A, Provenzano E, Wishart G, Pinder S, Watson P, Markowitz F, Murphy L, Ellis I, Purushotham A, Borresen-Dale AL, Brenton JD, Tavar   S, Caldas C, Aparicio S. The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature* 486:346-352, 2012.
8. Yin M, Lu M, Yao G, Tian H, Lian J, Liu L, Liang M, Wang Y, Sun F. Transactivation of microRNA-383 by steroidogenic factor-1 promotes estradiol release from mouse ovarian granulosa cells by targeting RBMS1. *Mol Endocrinol* 26:1129-1143, 2012.
9. Jiang J, Xie C, Liu Y, Shi Q, Chen Y. Up-regulation of miR-383-5p suppresses proliferation and enhances chemosensitivity in ovarian cancer cells by targeting TRIM27. *Biomed Pharmacother* 109:595-601, 2019.
10. Wang Z, Zhou Y, Hu X, Chen W, Lin X, Sun L, Xu X, Hong W, Wang T. RILP suppresses invasion of breast cancer cells by modulating the activity of RalA through interaction with RalGDS. *Cell Death Dis* 6:e1923, 2015.
11. Ward Y, Wang W, Woodhouse E, Linnoila I, Liotta L, Kelly K. Signal pathways which promote invasion and metastasis: Critical and distinct contributions of extracellular signal-regulated kinase and ral-specific guanine exchange factor pathways. *Mol Cell Biol* 21:5958-5969, 2001.
12. Bhattacharya M, Anborgh PH, Babwah AV, Dale LB, Dobransky T, Benovic JL, Feldman RD, Verdi JM, Rylett RJ, Ferguson SSG. Beta-arrestins regulate a ral-GDS ral effector pathway that mediates cytoskeletal reorganization. *Nat Cell Biol* 4:547-555, 2002.
13. Lee YH, Kim JH, Song GG. Genome-wide pathway analysis of breast cancer. *Tumour Biol* 35:7699-7705, 2014.
14. Conway K, Edmiston SN, May R, Kuan PF, Chu H, Bryant C, Tse CK, Swift-Scanlan T, Geradts J, Troester MA, Millikan RC. DNA methylation profiling in the carolina breast cancer study defines cancer subclasses differing in clinicopathologic characteristics and survival. *Breast Cancer Res* 16:45-6, 2014.
15. Ronneberg JA, Fleischer T, Solvang HK, Nordgard SH, Edvardsen H, Potapenko I, Nebdal D, Daviaud C, Gut I, Bukholm I, Naume B, Borresen-Dale AL, Tost J, Kristensen V. Methylation profiling with a panel of cancer related genes: Association with estrogen receptor, TP53 mutation status and expression subtypes in sporadic breast cancer. *Mol Oncol* 5:61-76, 2011.
16. Chang JW, Kuo WH, Lin CM, Chen WL, Chan SH, Chiu MF, Chang IS, Jiang SS, Tsai FY, Chen CH, Huang PH, Chang KJ, Lin KT, Lin SC, Wang MY, Uen YH, Tu CW, Hou MF, Tsai SF, Shen CY, Tung SL, Wang LH. Wild-type p53 upregulates an early onset breast cancer-associated gene GAS7 to suppress metastasis via GAS7-CYFIP1-mediated signaling pathway. *Oncogene* 37:4137-4150, 2018.
17. Gyorffy B, Lanczky A, Eklund AC, Denkert C, Budczies J, Li Q, Szallasi Z. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast Cancer Res Treat* 123:725-731, 2010.
18. Hung F, Chao CC-. Knockdown of growth-arrest-specific gene 7b (gas7b) using short-hairpin RNA desensitizes neuroblastoma cells to cisplatin: Implications for preventing apoptosis of neurons. *J Neurosci Res* 88:3578-3587, 2010.
19. Li D, Zhang B, Hu C. Oxaliplatin inhibits proliferation and migration of human hepatocellular carcinoma cells via GAS7C and the N-WASP/FAK/F-actin pathway. *Acta Biochim Biophys Sin (Shanghai)* 49:581-587, 2017.
20. Ping W, Gao Y, Fan X, Li W, Deng Y, Fu X. MiR-181a contributes gefitinib resistance in non-small cell lung cancer cells by targeting GAS7. *Biochem Biophys Res Commun* 495:2482-2489, 2018.
21. Michailidou K, Hall P, Gonzalez-Neira A, Ghoussaini M, Dennis J, Milne RL, Schmidt MK, Chang-Claude J, Bojesen SE, Bolla MK, Wang Q, Dicks E, Lee A, Turnbull C, Rahman N, Breast and Ovarian Cancer Susceptibility Collaboration, Fletcher O, Peto J, Gibson L, Dos Santos Silva I, Nevanlinna H, Muranen TA, Aittomaki K, Blomqvist C, Czene K, Irwanto A, Liu J, Waisfisz Q, Meijers-Heijboer H, Adank M, Hereditary Breast and Ovarian Cancer Research Group Netherlands, (HEBON), van der Luit, R. B., Hein R, Dahmen N, Beckman L, Meindl A, Schmutzler RK, Muller-Myhsok B, Lichtner P, Hopper JL, Southey MC, Makalic E, Schmidt DF, Uitterlinden AG, Hofman A, Hunter DJ, Chanock SJ, Vincent D, Baccot F, Tessier DC, Canisius S, Wessels LF, Haiman CA, Shah M, Luben R, Brown J, Luccarini C, Schoof N, Humphreys K, Li J, Nordestgaard BG, Nielsen SF, Flyger H, Couch FJ, Wang X, Vachon C, Stevens KN, Lambrechts D, Moisse M, Paridaens R, Christiaens MR, Rudolph A, Nickels S, Flesch-Janys D, Johnson N, Aitken Z, Aaltonen K, Heikinen T, Broeks A, Veer LJ, van der Schoot, C. E., Guenel P, Truong T, Laurent-Puig P, Menegaux F, Marme F, Schneeweiss A, Sohn C, Burwinkel B, Zamora MP, Perez JL, Pita G, Alonso MR, Cox A, Brock IW, Cross SS, Reed MW, Sawyer EJ, Tomlinson I, Kerin MJ, Miller N, Henderson BE, Schumacher F, Le Marchand L, Andrulis IL, Knight JA, Glendon G, Mulligan AM, kConFab Investigators, stralian Ovarian Cancer Study Group, Lindblom A, Margolin S, Hooning MJ, Hollestelle A, van den Ouweland, A. M., Jager A, Bui QM, Stone J, Dite GS, Apicella C, Tsimiklis H, Giles GG, Severi G, Baglietto L, Fasching PA, Haeberle L, Ekici AB, Beckmann MW, Brenner H, Muller H, Arndt V, Stegmaier C, Swerdlow A, Ashworth A, Orr N, Jones M, Figueroa J, Lissowska

- J, Brinton L, Goldberg MS, Labreche F, Dumont M, Winqvist R, Pylkas K, Jukkola-Vuorinen A, Grip M, Brauch H, Hamann U, Bruning T, GENICA (Gene Environment Interaction and Breast Cancer in Germany) Network, Radice P, Peterlongo P, Manoukian S, Bonanni B, Devilee P, Tollenaar RA, Seynaeve C, van Asperen CJ, Jakubowska A, Lubinski J, Jaworska K, Durda K, Mannermaa A, Kataja V, Kosma VM, Hartikainen JM, Bogdanova NV, Antonenkova NN, Dork T, Kristensen VN, Anton-Culver H, Slager S, Toland AE, Edge S, Fostira F, Kang D, Yoo KY, Noh DY, Matsuo K, Ito H, Iwata H, Sueta A, Wu AH, Tseng CC, Van Den Berg D, Stram DO, Shu XO, Lu W, Gao YT, Cai H, Teo SH, Yip CH, Phuah SY, Cornes BK, Hartman M, Miao H, Lim WY, Sng JH, Muir K, Lophatananon A, Stewart-Brown S, Siriwanarangsana P, Shen CY, Hsiung CN, Wu PE, Ding SL, Sangrajrang S, Gaborieau V, Brennan P, McKay J, Blot WJ, Signorello LB, Cai Q, Zheng W, Deming-Halverson S, Shrubsole M, Long J, Simard J, Garcia-Closas M, Pharoah PD, Chenevix-Trench G, Dunning AM, Benitez J, Easton DF. Large-scale genotyping identifies 41 new loci associated with breast cancer risk. *Nat Genet* 45:353-361, 2013.
22. Yuan J, Zhang N, Zhu H, Liu J, Xing H, Ma F, Yang M. CHST9 rs1436904 genetic variant contributes to prognosis of triple-negative breast cancer. *Sci Rep* 7:1180-6, 2017.
23. Schabla NM, Mondal K, Swanson PC. DCAF1 (VprBP): Emerging physiological roles for a unique dual-service E3 ubiquitin ligase substrate receptor. *J Mol Cell Biol* , 2018.
24. Britschgi A, Duss S, Kim S, Couto JP, Brinkhaus H, Koren S, De Silva D, Mertz KD, Kaup D, Varga Z, Voshol H, Vissieres A, Leroy C, Roloff T, Stadler MB, Scheel CH, Miraglia LJ, Orth AP, Bonamy GMC, Reddy VA, Bentires-Alj M. The hippo kinases LATS1 and 2 control human breast cell fate via crosstalk with ERα. *Nature* 541:541-545, 2017.
25. Lee JM, Lee JS, Kim H, Kim K, Park H, Kim J, Lee SH, Kim IS, Kim J, Lee M, Chung CH, Seo S, Yoon J, Ko E, Noh D, Kim KI, Kim KK, Baek SH. EZH2 generates a methyl degron that is recognized by the DCAF1/DBP1/CUL4 E3 ubiquitin ligase complex. *Mol Cell* 48:572-586, 2012.
26. Guo Z, Kong Q, Liu C, Zhang S, Zou L, Yan F, Whitmire JK, Xiong Y, Chen X, Wan YY. DCAF1 controls T-cell function via p53-dependent and -independent mechanisms. *Nat Commun* 7:10307, 2016.
27. Hossain D, Ferreira Barbosa JA, Cohen ÉA, Tsang WY. HIV-1 vpr hijacks EDD-DYRK2-DBP1DCAF1 to disrupt centrosome homeostasis. *J Biol Chem* 293:9448-9460, 2018.
28. Zhang W, Wu M, Chong Q, Zhang M, Zhang X, Hu L, Zhong Y, Qian P, Kong X, Tan S, Li G, Ding K, Lobie PE, Zhu T. Loss of estrogen-regulated MIR135A1 at 3p21.1 promotes tamoxifen resistance in breast cancer. *Cancer Res* 78:4915-4928, 2018.
29. Yan L, Chen Z, Li-Li n, Chen J, Wei W, Mo X, Qin Y, Lin Y, Chen J. miR-135a promotes gastric cancer progression and resistance to oxaliplatin. *Oncotarget* 7:70699-70714, 2016.
30. Zhang T, Wang N. miR-135a confers resistance to gefitinib in non-small cell lung cancer cells by upregulation of RAC1. *Oncol Res* 26:1191-1200, 2018.
31. Maitra A, Wistuba II, Washington C, Virmani AK, Ashfaq R, Milchgrub S, Gazdar AF, Minna JD. High-resolution chromosome 3p allelotyping of breast carcinomas and precursor lesions demonstrates frequent loss of heterozygosity and a discontinuous pattern of allele loss. *Am J Pathol* 159:119-130, 2001.
32. Martinez A, Walker RA, Shaw JA, Dearing SJ, Maher ER, Latif F. Chromosome 3p allele loss in early invasive breast cancer: Detailed mapping and association with clinicopathological features. *MP, Mol Pathol* 54:300-306, 2001.
33. Zhang P, Xiao Z, Wang S, Zhang M, Wei Y, Hang Q, Kim J, Yao F, Rodriguez-Aguayo C, Ton BN, Lee M, Wang Y, Zhou Z, Zeng L, Hu X, Lawhon SE, Siverly AN, Su X, Li J, Xie X, Cheng X, Liu L, Chang H, Chiang S, Lopez-Berestein G, Sood AK, Chen J, You MJ, Sun S, Liang H, Huang Y, Yang X, Sun D, Sun Y, Hung M, Ma L. ZRANB1 is an EZH2 deubiquitinase and a potential therapeutic target in breast cancer. *Cell Rep* 23:823-837, 2018.
34. Bai SW, Herrera-Abreu MT, Rohn JL, Racine V, Tajadura V, Suryavanshi N, Bechtel S, Wiemann S, Baum B, Ridley AJ. Identification and characterization of a set of conserved and new regulators of cytoskeletal organization, cell morphology and migration. *BMC Biol* 9:54, 2011.
35. Jin J, Xie X, Xiao Y, Hu H, Zou Q, Cheng X, Sun S. Epigenetic regulation of the expression of IL12 and IL23 and autoimmune inflammation by the deubiquitinase TRABID. *Nat Immunol* 17:259-268, 2016.
36. Bergman LM, Birts CN, Darley M, Gabrielli B, Blaydes JP. CtBPs promote cell survival through the maintenance of mitotic fidelity. *Mol Cell Biol* 29:4539-4551, 2009.
37. Kim TW, Kang B, Jang H, Kwak S, Shin J, Kim H, Lee S, Lee S, Lee J, Kim J, Kim S, Cho E, Kim JH, Park KS, Che J, Han DW, Kang MJ, Yi EC, Youn H. Ctbp2 modulates NuRD-mediated deacetylation of H3K27 and facilitates PRC2-mediated H3K27me3 in active embryonic stem cell genes during exit from pluripotency. *Stem Cells* 33:2442-2455, 2015.
38. Yang X, Sun Y, Li H, Shao Y, Zhao D, Yu W, Fu J. C-terminal binding protein-2 promotes cell proliferation and migration in breast cancer via suppression of p16INK4A. *Oncotarget* 8:26154-26168, 2017.
39. Cui TX, Kryczek I, Zhao L, Zhao E, Kuick R, Roh MH, Vatan L, Szeliga W, Mao Y, Thomas DG, Kotarski J, Tarkowski R, Wicha M, Cho K, Giordano T, Liu R, Zou W. Myeloid-derived suppressor cells enhance stemness of cancer cells by inducing microRNA101 and suppressing the corepressor CtBP2. *Immunity* 39:611-621, 2013.
40. Birts CN, Harding R, Soosaipillai G, Halder T, Azim-Araghi A, Darley M, Cutress RI, Bateman AC, Blaydes JP. Expression of CtBP family protein isoforms in breast cancer and their role in chemoresistance. *Biol Cell* 103:1-19, 2010.
41. Li L, Wan S, Tao K, Wang G, Zhao E. KLRG1 restricts memory T cell antitumor immunity. *Oncotarget* 7:61670-61678, 2016.
42. May T, Yang J, Shoni M, Liu S, He H, Gali R, Ng S, Crum C, Berkowitz RS, Ng S. BRCA1 expression is epigenetically repressed in sporadic ovarian cancer cells by overexpression of C-terminal binding protein 2. *Neoplasia* 15:600-608, 2013.
43. Wang DP, Gu LL, Xue Q, Chen H, Mao GX. CtBP2 promotes proliferation and reduces drug sensitivity in non-small cell lung cancer via the wnt/β-catenin pathway. *Neoplasia* 65:888-897, 2018.



44. Shi H, Mao Y, Ju Q, Wu Y, Bai W, Wang P, Zhang Y, Jiang M. C-terminal binding protein 2 mediates cisplatin chemoresistance in esophageal cancer cells via the inhibition of apoptosis. *Int J Oncol* 53:167-176, 2018.
45. Thomas G, Jacobs KB, Kraft P, Yeager M, Wacholder S, Cox DG, Hankinson SE, Hutchinson A, Wang Z, Yu K, Chatterjee N, Garcia-Closas M, Gonzalez-Bosquet J, Prokunina-Olsson L, Orr N, Willett WC, Colditz GA, Ziegler RG, Berg CD, Buys SS, McCarty CA, Feigelson HS, Calle EE, Thun MJ, Diver R, Prentice R, Jackson R, Kooperberg C, Chlebowski R, Lissowska J, Peplonska B, Brinton LA, Sigurdson A, Doody M, Bhatti P, Alexander BH, Buring J, Lee IM, Vatten LJ, Hveem K, Kumle M, Hayes RB, Tucker M, Gerhard DS, Fraumeni JF, Jr, Hoover RN, Chanock SJ, Hunter DJ. A multistage genome-wide association study in breast cancer identifies two new risk alleles at 1p11.2 and 14q24.1 (RAD51L1). *Nat Genet* 41:579-584, 2009.
46. Suwaki N, Klare K, Tarsounas M. RAD51 paralogs: Roles in DNA damage signalling, recombinational repair and tumorigenesis. *Semin Cell Dev Biol* 22:898-905, 2011.
47. Zhu B, Mukherjee A, Machiela MJ, Song L, Hua X, Shi J, Garcia-Closas M, Chanock SJ, Chatterjee N. An investigation of the association of genetic susceptibility risk with somatic mutation burden in breast cancer. *Br J Cancer* 115:752-760, 2016.
48. Takada M, Nagai S, Haruta M, Sugino RP, Tozuka K, Takei H, Ohkubo F, Inoue K, Kurosumi M, Miyazaki M, Sato-Otsubo A, Sato Y, Ogawa S, Kaneko Y. BRCA1 alterations with additional defects in DNA damage response genes may confer chemoresistance to BRCA-like breast cancers treated with neoadjuvant chemotherapy. *Genes Chromosomes Cancer* 56:405-420, 2017.
49. Bhardwaj A, Singh H, Rajapakshe K, Tachibana K, Ganesan N, Pan Y, Gunaratne PH, Coarfa C, Bedrosian I. Regulation of miRNA-29c and its downstream pathways in preneoplastic progression of triple-negative breast cancer. *Oncotarget* 8:19645-19660, 2017.
50. Kadota M, Sato M, Duncan B, Ooshima A, Yang HH, Diaz-Meyer N, Gere S, Kageyama S, Fukuoka J, Nagata T, Tsukada K, Dunn BK, Wakefield LM, Lee MP. Identification of novel gene amplifications in breast cancer and coexistence of gene amplification with an activating mutation of PIK3CA. *Cancer Res* 69:7357-7365, 2009.
51. Shimoda M, Hori A, Wands JR, Tsunashima R, Naoi Y, Miyake T, Tanei T, Kagara N, Shimazu K, Kim SJ, Noguchi S. Endocrine sensitivity of estrogen receptor-positive breast cancer is negatively correlated with aspartate- $\beta$ -hydroxylase expression. *Cancer Sci* 108:2454-2461, 2017.
52. Tomimaru Y, Mishra S, Safran H, Charpentier KP, Martin W, De Groot AS, Gregory SH, Wands JR. Aspartate- $\beta$ -hydroxylase induces epitope-specific T cell responses in hepatocellular carcinoma. *Vaccine* 33:1256-1266, 2015.
53. Shimoda M, Tomimaru Y, Charpentier KP, Safran H, Carlson RI, Wands J. Tumor progression-related transmembrane protein aspartate- $\beta$ -hydroxylase is a target for immunotherapy of hepatocellular carcinoma. *J Hepatol* 56:1129-1135, 2012.
54. Kouros-Mehr H, Slorach EM, Sternlicht MD, Werb Z. GATA-3 maintains the differentiation of the luminal cell fate in the mammary gland. *Cell* 127:1041-1055, 2006.
55. Asselin-Labat M, Sutherland KD, Barker H, Thomas R, Shackleton M, Forrest NC, Hartley L, Robb L, Grosveld FG, van der Wees J, Lindeman GJ, Visvader JE. Gata-3 is an essential regulator of mammary-gland morphogenesis and luminal-cell differentiation. *Nat Cell Biol* 9:201-209, 2007.
56. Theodorou V, Stark R, Menon S, Carroll JS. GATA3 acts upstream of FOXA1 in mediating ESR1 binding by shaping enhancer accessibility. *Genome Res* 23:12-22, 2013.
57. Shahi P, Wang C-, Chou J, Hagerling C, Gonzalez Velozo H, Ruderisch A, Yu Y, Lai M-, Werb Z. GATA3 targets semaphorin 3B in mammary epithelial cells to suppress breast cancer progression and metastasis. *Oncogene* 36:5567-5575, 2017.
58. Mehra R, Varambally S, Ding L, Shen R, Sabel MS, Ghosh D, Chinnaiyan AM, Kleer CG. Identification of GATA3 as a breast cancer prognostic marker by global gene expression meta-analysis. *Cancer Res* 65:11259-11264, 2005.
59. Yoon, Nam K., BS|Maresh, Erin L., BS|Shen, Dejun, MD|Elshimali, Yahya, MD|Apple, Sophia, MD|Horvath, Steve, PhD|Mah, Vei, MD|Bose, Shikha, MD|Chia, David, PhD|Chang, Helena R., MD|Goodglick, Lee, PhD. Higher levels of GATA3 predict better survival in women with breast cancer. *Human Pathology* 41:1794-1801, 2010.
60. Albergaria A, Paredes J, Sousa B, Milanezi F, Carneiro V, Bastos J, Costa S, Vieira D, Lopes N, Lam EW, Lunet N, Schmitt F. Expression of FOXA1 and GATA-3 in breast cancer: The prognostic significance in hormone receptor-negative tumours. *Breast Cancer Res* 11:R40, 2009.
61. Tominaga N, Naoi Y, Shimazu K, Nakayama T, Maruyama N, Shimomura A, Kim SJ, Tamaki Y, Noguchi S. Clinicopathological analysis of GATA3-positive breast cancers with special reference to response to neoadjuvant chemotherapy. *Ann Oncol* 23:3051-3057, 2012.
62. Gulbahce, H. Evin, MD|Sweeney, Carol, PhD|Surowiecka, Maria, MD|Knapp, Dennis, BS|Varghese, Linda, MD|Blair, Cindy K., PhD. Significance of GATA-3 expression in outcomes of patients with breast cancer who received systemic chemotherapy and/or hormonal therapy and clinicopathologic features of GATA-3-positive tumors. *Human Pathology* 44:2427-2431, 2013.
63. Ciocca, Vincenzo, DO|Daskalakis, Constantine, ScD|Ciocca, Robin M., DO|Ruiz-Orrico, Alejandra, MD|Palazzo, Juan P., MD. The significance of GATA3 expression in breast cancer: A 10-year follow-up study. *Human Pathology* 40:489-495, 2009.
64. Jenssen T-, Kuo WP, Stokke T, Hovig E. Associations between gene expressions in breast cancer and patient survival. *Hum Genet* 111:411-420, 2002.
65. Liu J, Prager-van der Smissen, Wendy J. C., Look MP, Sieuwerts AM, Smid M, Meijer-van Gelder ME, Foekens JA, Hollestelle A, Martens JWM. GATA3 mRNA expression, but not mutation, associates with longer progression-free survival in ER-positive breast cancer patients treated with first-line tamoxifen for recurrent disease. *Cancer Lett* 376:104-109, 2016.
66. Bollet MA, Savignoni A, De Koning L, Tran-Perennou C, Barbaroux C, Degeorges A, Sigal-Zafrani B, Almouzni G, Cottu P, Salmon R, Servant N, Fourquet A, de Cremoux P. Tumor aromatase expression as a

- prognostic factor for local control in young breast cancer patients after breast-conserving treatment. *Breast Cancer Res* 11:R54, 2009.
67. Shafer MER, Nguyen AHT, Tremblay M, Viala S, Béland M, Bertos NR, Park M, Bouchard M. Lineage specification from prostate progenitor cells requires Gata3-dependent mitotic spindle orientation. *Stem Cell Reports* 8:1018-1031, 2017.
  68. Wan YY. GATA3: A master of many trades in immune regulation. *Trends Immunol* 35:233-242, 2014.
  69. Bian C, Chen Q, Yu X. The zinc finger proteins ZNF644 and WIZ regulate the G9a/GLP complex for gene repression. *Elife* 4, 2015.
  70. Mozzetta C, Pontis J, Fritsch L, Robin P, Portoso M, Proux C, Margueron R, Ait-Si-Ali S. The histone H3 lysine 9 methyltransferases G9a and GLP regulate polycomb repressive complex 2-mediated gene silencing. *Mol Cell* 53:277-289, 2014.
  71. Dungrawala H, Rose KL, Bhat KP, Mohni KN, Glick GG, Couch FB, Cortez D. The replication checkpoint prevents two types of fork collapse without regulating replisome stability. *Mol Cell* 59:998-1010, 2015.
  72. Maiato H, Fairley EAL, Rieder CL, Swedlow JR, Sunkel CE, Earnshaw WC. Human CLASP1 is an outer kinetochore component that regulates spindle microtubule dynamics. *Cell* 113:891-904, 2003.
  73. Pereira AL, Pereira AJ, Maia ARR, Drabek K, Sayas CL, Hergert PJ, Lince-Faria M, Matos I, Duque C, Stepanova T, Rieder CL, Earnshaw WC, Galjart N, Maiato H. Mammalian CLASP1 and CLASP2 cooperate to ensure mitotic fidelity by regulating spindle and kinetochore function. *Mol Biol Cell* 17:4526-4542, 2006.
  74. Bouchet BP, Noordstra I, van Amersfoort M, Katrukha EA, Ammon Y, Ter Hoeve ND, Hodgson L, Dogterom M, Derksen PWB, Akhmanova A. Mesenchymal cell invasion requires cooperative regulation of persistent microtubule growth by SLAIN2 and CLASP1. *Dev Cell* 39:708-723, 2016.
  75. Lin T, Su C, Wu P, Lai T, Pan W, Jan Y, Chang Y, Yeh C, Chen C, Ger L, Chang H, Yang C, Huang M, Liu Y, Lin Y, Shyy JY-, Tsai M, Hsiao M. The nucleolar protein NIFK promotes cancer progression via CK1 $\alpha$ / $\beta$ -catenin in metastasis and ki-67-dependent cell proliferation. *Elife* 5, 2016.
  76. Pan W, Tsai H, Wang S, Hsiao M, Wu P, Tsai M. The RNA recognition motif of NIFK is required for rRNA maturation during cell cycle progression. *RNA Biol* 12:255-267, 2015.
  77. Zhou Y, Zhao W, Mao L, Wang Y, Xia L, Cao M, Shen J, Chen J. Long non-coding RNA NIFK-AS1 inhibits M2 polarization of macrophages in endometrial cancer through targeting miR-146a. *Int J Biochem Cell Biol* 104:25-33, 2018.
  78. Kotarba G, Krzywinska E, Grabowska AI, Taracha A, Wilanowski T. TFCEP2/TFCEP2L1/UBP1 transcription factors in cancer. *Cancer Lett* 420:72-79, 2018.
  79. Liu Y, Shao Z, Yuan G. Prediction of polycomb target genes in mouse embryonic stem cells. *Genomics* 96:17-26, 2010.
  80. Wang H, Li M, Tao X, Qian Y, Chen L, Tao G. ARHGEF39 promotes gastric cancer cell proliferation and migration via akt signaling pathway. *Mol Cell Biochem* 440:33-42, 2018.
  81. Zhou H, Cai L, Zhang X, Li A, Miao Y, Li Q, Qiu X, Wang E. ARHGEF39 promotes tumor progression via activation of Rac1/P38 MAPK/ATF2 signaling and predicts poor prognosis in non-small cell lung cancer patients. *Lab Invest* 98:670-681, 2018.
  82. Zhang J, Zhang J, Xu S, Zhang X, Wang P, Wu H, Xia B, Zhang G, Lei B, Wan L, Zhang D, Pang D. Hypoxia-induced TPM2 methylation is associated with chemoresistance and poor prognosis in breast cancer. *Cell Physiol Biochem* 45:692-705, 2018.
  83. Groth-Pedersen L, Aits S, Corcelle-Termeau E, Petersen NHT, Nylandsted J, Jäättelä M. Identification of cytoskeleton-associated proteins essential for lysosomal stability and survival of human cancer cells. *PLoS ONE* 7:e45381, 2012.
  84. Astudillo L, Therville N, Colacios C, Ségui B, Andrieu-Abadie N, Levade T. Glucosylceramidases and malignancies in mammals. *Biochimie* 125:267-280, 2016.
  85. Raju D, Schonauer S, Hamzeh H, Flynn KC, Bradke F, Vom Dorp K, Dörmann P, Yildiz Y, Trötschel C, Poetsch A, Breiden B, Sandhoff K, Körschen HG, Wachten D. Accumulation of glucosylceramide in the absence of the beta-glucosidase GBA2 alters cytoskeletal dynamics. *PLoS Genet* 11:e1005063, 2015.
  86. Duan B, Cui J, Sun S, Zheng J, Zhang Y, Ye B, Chen Y, Deng W, Du J, Zhu Y, Chen Y, Gu L. EGF-stimulated activation of Rab35 regulates RUSC2-GIT2 complex formation to stabilize GIT2 during directional lung cancer cell migration. *Cancer Lett* 379:70-83, 2016.
  87. Zheng M, Xing C, Xiao H, Ma N, Wang X, Han G, Chen G, Hou C, Shen B, Li Y, Wang R. Interaction of CD5 and CD72 is involved in regulatory T and B cell homeostasis. *Immunol Invest* 43:705-716, 2014.
  88. Jiang X, Björkström NK, Melum E. Intact CD100-CD72 interaction necessary for TCR-induced T cell proliferation. *Front Immunol* 8:765, 2017.

## Supplementary note

### Acknowledgements

We thank all the individuals who took part in these studies and all the researchers, clinicians, technicians and administrative staff who have enabled this work to be carried out; Sue Healey, in particular taking on the task of mutation classification with the late Olga Sinilnikova; Maggie Angelakos, Judi Maskiell, Gillian Dite, Helen Tsimiklis; members and participants in the New York site of the Breast Cancer Family Registry; members and participants in the Ontario Familial Breast Cancer Registry; Vilius Rudaitis and Laimonas Griškevičius; Drs Janis Eglitis, Anna Krilova and Aivars Stengrevics; Yuan Chun Ding and Linda Steele for their work in participant enrollment and biospecimen and data management; Bent Ejlersen and Anne-Marie Gerdes for the recruitment and genetic counseling of participants; Alicia Barroso, Rosario Alonso and Guillermo Pita; Manoukian Siranoush, Bernard Peissel, Cristina Zanzottera, Milena Mariani, Daniela Zaffaroni, Bernardo Bonanni, Monica Barile, Irene Feroce, Mariarosaria Calvello, Alessandra Viel, Riccardo Dolcetti, Laura Ottini, Giuseppe Giannini, Laura Papi, Gabriele Lorenzo Capone, Liliana Varesco, Viviana Gismondi, Maria Grazia Tibiletti, Daniela Furlan, Antonella Savarese, Aline Martayan, Stefania Tommasi, Brunella Pilato; the personnel of the Cogentech Cancer Genetic Test Laboratory, Milan, Italy. Ms. JoEllen Weaver and Dr. Betsy Bove; Marta Santamariña, Ana Blanco, Miguel Aguado, Uxía Esperón and Belinda Rodríguez; IFE - Leipzig Research Centre for Civilization Diseases (Markus Loeffler, Joachim Thiery, Matthias Nüchter, Ronny Baber); We thank all participants, clinicians, family doctors, researchers, and technicians for their contributions and commitment to the DKFZ study and the collaborating groups in Lahore, Pakistan (Muhammad U. Rashid, Noor Muhammad, Sidra Gull, Seerat Bajwa, Faiz Ali Khan, Humaira Naeemi, Saima Faisal, Asif Loya, Mohammed Aasim Yusuf) and Bogota, Colombia (Diana Torres, Ignacio Briceno, Fabian Gil). Genetic Modifiers of Cancer Risk in BRCA1/2 Mutation Carriers (GEMO) study is a study from the National Cancer Genetics Network UNICANCER Genetic Group, France. We wish to pay a tribute to Olga M. Sinilnikova, who with Dominique Stoppa-Lyonnet initiated and coordinated GEMO until she sadly passed away on the 30th June 2014. The team in Lyon (Olga Sinilnikova, Mélanie Léone, Laure Barjhoux, Carole Verny-Pierre, Sylvie Mazoyer, Francesca Damiola, Valérie Sornin) managed the GEMO samples until the biological resource centre was transferred to Paris in December 2015 (Noura Mebirouk, Fabienne Lesueur, Dominique Stoppa-Lyonnet). We want to thank all the GEMO collaborating groups for their contribution to this study: Coordinating Centre, Service de Génétique, Institut Curie, Paris, France: Muriel Belotti, Ophélie Bertrand, Anne-Marie Birot, Bruno Buecher, Sandrine Caputo, Anaïs Dupré, Emmanuelle Fourme, Marion Gauthier-Villars, Lisa Golmard, Claude Houdayer, Marine Le Mentec, Virginie Moncoutier, Antoine de Pauw, Claire Saule, Dominique Stoppa-Lyonnet, and Inserm U900, Institut Curie, Paris, France: Fabienne Lesueur, Noura Mebirouk. Contributing Centres : Unité Mixte de Génétique Constitutionnelle des Cancers Fréquents, Hospices Civils de Lyon - Centre Léon Bérard, Lyon, France: Nadia Boutry-Kryza, Alain Calender, Sophie Giraud, Mélanie Léone. Institut Gustave Roussy, Villejuif, France: Brigitte Bressac-de-Paillerets, Olivier Caron, Marine Guillaud-Bataille. Centre Jean Perrin, Clermont-Ferrand, France: Yves-Jean Bignon, Nancy Uhrhammer. Centre Léon Bérard, Lyon, France: Valérie Bonadona, Christine Lasset. Centre François Baclesse, Caen, France: Pascaline Berthet, Laurent Castera, Dominique Vaur. Institut Paoli Calmettes, Marseille, France: Violaine Bourdon, Catherine Noguès, Tetsuro Noguchi, Cornel Popovici, Audrey Remenieras, Hagay Sobol. CHU Arnaud-de-Villeneuve, Montpellier, France: Isabelle Coupier, Pascal Pujol. Centre Oscar Lambret, Lille, France: Claude Adenis, Aurélie Dumont,

Françoise Révillion. Centre Paul Strauss, Strasbourg, France: Danièle Muller. Institut Bergonié, Bordeaux, France: Emmanuelle Barouk-Simonet, Françoise Bonnet, Virginie Bubien, Michel Longy, Nicolas Sevenet, Institut Claudius Regaud, Toulouse, France: Laurence Gladieff, Rosine Guimbaud, Viviane Feillel, Christine Toulas. CHU Grenoble, France: Hélène Dreyfus, Christine Dominique Leroux, Magalie Peysselon, Rebischung. CHU Dijon, France: Amandine Baurand, Geoffrey Bertolone, Fanny Coron, Laurence Faivre, Caroline Jacquot, Sarab Lizard. CHU St-Etienne, France: Caroline Kientz, Marine Lebrun, Fabienne Prieur. Hôtel Dieu Centre Hospitalier, Chambéry, France: Sandra Fert Ferrer. Centre Antoine Lacassagne, Nice, France: Véronique Mari. CHU Limoges, France: Laurence Vénat-Bouvet. CHU Nantes, France: Stéphane Bézieau, Capucine Delnatte. CHU Bretonneau, Tours and Centre Hospitalier de Bourges France: Isabelle Mortemousque. Groupe Hospitalier Pitié-Salpêtrière, Paris, France: Chrystelle Colas, Florence Coulet, Florent Soubrier, Mathilde Warcoin. CHU Vandoeuvre-les-Nancy, France: Myriam Bronner, Johanna Sokolowska. CHU Besançon, France: Marie-Agnès Collonge-Rame, Alexandre Damette. CHU Poitiers, Centre Hospitalier d'Angoulême and Centre Hospitalier de Niort, France: Paul Gesta. Centre Hospitalier de La Rochelle : Hakima Lallaoui. CHU Nîmes Carêmeau, France : Jean Chiesa. CHI Poissy, France: Denise Molina-Gomes. CHU Angers, France : Olivier Ingster; Ilse Coene en Brecht Crombez; Ilse Coene and Brecht Crombez; Alicia Tosar and Paula Diaque; Irja Erkkilä and Virpi Palola; HEBON thanks the study participants and the registration teams of IKNL and PALGA for part of the data collection; Hong Kong Sanatorium and Hospital; the Hungarian Breast and Ovarian Cancer Study Group members (Janos Papp, Aniko Bozsik, Timea Pocza, Zoltan Matrai, Miklos Kasler, Judit Franko, Maria Balogh, Gabriella Domokos, Judit Ferenczi, Department of Molecular Genetics, National Institute of Oncology, Budapest, Hungary) and the clinicians and patients for their contributions to this study; the Oncogenetics Group (VHIO) and the High Risk and Cancer Prevention Unit of the University Hospital Vall d'Hebron, and the Cellex Foundation for providing research facilities and equipment; the ICO Hereditary Cancer Program team led by Dr. Gabriel Capella; the ICO Hereditary Cancer Program team led by Dr. Gabriel Capella; Dr Martine Dumont for sample management and skillful assistance; Ana Peixoto, Catarina Santos and Pedro Pinto; members of the Center of Molecular Diagnosis, Oncogenetics Department and Molecular Oncology Research Center of Barretos Cancer Hospital; all the kConFab research nurses and staff, the heads and staff of the Family Cancer Clinics, and the Clinical Follow Up Study (which has received funding from the NHMRC, the National Breast Cancer Foundation, Cancer Australia, and the National Institute of Health (USA)) for their contributions to this resource, and the many families who contribute to kConFab; the KOBRA Study Group; Csilla Szabo (National Human Genome Research Institute, National Institutes of Health, Bethesda, MD, USA); Lenka Foretova and Eva Machackova (Department of Cancer Epidemiology and Genetics, Masaryk Memorial Cancer Institute and MF MU, Brno, Czech Republic); and Michal Zikan, Petr Pohlreich and Zdenek Kleibl (Oncogynecologic Center and Department of Biochemistry and Experimental Oncology, First Faculty of Medicine, Charles University, Prague, Czech Republic); Anne Lincoln, Lauren Jacobs; the NICCC National Familial Cancer Consultation Service team led by Sara Dishon, the lab team led by Dr. Flavio Lejbkowitz, and the research field operations team led by Dr. Mila Pinchev; the investigators of the Australia New Zealand NRG Oncology group; members and participants in the Ontario Cancer Genetics Network; Leigha Senter, Kevin Sweet, Caroline Craven, Julia Cooper, and Michelle O'Connor; Yip Cheng Har, Nur Aishah Mohd Taib, Phuah Sze Yee, Norhashimah Hassan and all the research nurses, research assistants and doctors involved in the MyBrCa Study for assistance in patient recruitment, data collection and sample preparation, Philip Iau, Sng Jen-Hwei and Sharifah Nor Akmal for contributing samples from the Singapore

Breast Cancer Study and the HUKM-HKL Study respectively; the Meirav Comprehensive breast cancer center team at the Sheba Medical Center; Christina Selkirk; Cecilia Zvocec, Qun Niu; Joyce Seldon and Lorna Kwan; Dr. Robert Nussbaum, Beth Crawford, Kate Loranger, Julie Mak, Nicola Stewart, Robin Lee, Amie Blanco and Peggy Conrad and Salina Chan; Simon Gayther, Susan Ramus, Paul Pharoah, Carole Pye, Patricia Harrington and Eva Wozniak; Geoffrey Lindeman, Marion Harris, Martin Delatycki, Sarah Sawyer, Rebecca Driessen, and Ella Thompson for performing all DNA amplification.

The CIMBA data management and data analysis were supported by Cancer Research – UK grants C12292/A20861, C12292/A11174. ACA is a Cancer Research -UK Senior Cancer Research Fellow. GCT and ABS are NHMRC Research Fellows. Genotyping for the OncoArray was funded by the government of Canada through Genome Canada and the Canadian Institutes of Health Research (GPH-129344), the Ministère de l'Économie, de la Science et de l'Innovation du Québec through Génome Québec, the Quebec Breast Cancer Foundation for the PERSPECTIVE project, the US National Institutes of Health (NIH) (1 U19 CA 148065 for the Discovery, Biology and Risk of Inherited Variants in Breast Cancer (DRIVE) project and X01HG007492 to the Center for Inherited Disease Research (CIDR) under contract HHSN268201200008I), Cancer Research UK (C1287/A16563), the Odense University Hospital Research Foundation (Denmark), the National R&D Program for Cancer Control–Ministry of Health and Welfare (Republic of Korea) (1420190), the Italian Association for Cancer Research (AIRC; IG16933), the Breast Cancer Research Foundation, the National Health and Medical Research Council (Australia) and German Cancer Aid (110837).

BCFR-AU / BCFR-NC: UM1 CA164920 from the U.S. National Cancer Institute. The content of this manuscript does not necessarily reflect the views or policies of the National Cancer Institute or any of the collaborating centers in the Breast Cancer Family Registry (BCFR), nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government or the BCFR.

BCFR-ON: The Ontario Familial Breast Cancer Registry was supported by grant U01CA164920 from the U.S. National Cancer Institute of the National Institutes of Health. The content of this manuscript does not necessarily reflect the views or policies of the National Cancer Institute or any of the collaborating centers in the Breast Cancer Family Registry (BCFR), nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government or the BCFR.

COH: Research reported in this publication was supported by the National Cancer Institute of the National Institutes of Health under grant number R25CA112486, and RC4CA153828 (PI: J. Weitzel) from the National Cancer Institute and the Office of the Director, National Institutes of Health. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

DEMOKRITOS: European Union (European Social Fund – ESF) and Greek national funds through the Operational Program "Education and Lifelong Learning" of the National Strategic Reference Framework (NSRF) - Research Funding Program of the General Secretariat for Research & Technology: SYN11\_10\_19 NBCA. Investing in knowledge society through the European Social Fund.

EMBRACE: Cancer Research UK Grants C1287/A10118 and C1287/A11990. D. Gareth Evans and Fiona Laloo are supported by an NIHR grant to the Biomedical Research Centre, Manchester. The Investigators at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust are supported by an NIHR grant to the Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust. Ros Eeles and Elizabeth Bancroft are supported by Cancer Research UK Grant

C5047/A8385. Ros Eeles is also supported by NIHR support to the Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust.

GC-HBOC: German Cancer Aid (grant no 110837, Rita K. Schmutzler) and the European Regional Development Fund and Free State of Saxony, Germany (LIFE - Leipzig Research Centre for Civilization Diseases, project numbers 713-241202, 713-241202, 14505/2470, 14575/2470).

HEBCS: Helsinki University Hospital Research Fund, Academy of Finland (266528), the Finnish Cancer Society and the Sigrid Juselius Foundation.

HEBON: the Dutch Cancer Society grants NKI1998-1854, NKI2004-3088, NKI2007-3756, the Netherlands Organization of Scientific Research grant NWO 91109024, the Pink Ribbon grants 110005 and 2014-187.WO76, the BBMRI grant NWO 184.021.007/CP46 and the Transcan grant JTC 2012 Cancer 12-054.

HEBON thanks the registration teams of Dutch Cancer Registry (IKNL; S. Siesling, J. Verloop) and the Dutch Pathology database (PALGA; L. Overbeek) for part of the data collection.

ICO: The authors would like to particularly acknowledge the support of the Asociación Española Contra el Cáncer (AECC), the Instituto de Salud Carlos III (organismo adscrito al Ministerio de Economía y Competitividad) and “Fondo Europeo de Desarrollo Regional (FEDER), una manera de hacer Europa” (PI10/01422, PI13/00285, PIE13/00022, PI15/00854, PI16/00563 and CIBERONC) and the Institut Català de la Salut and Autonomous Government of Catalonia (2009SGR290, 2014SGR338 and PERIS Project MedPerCan).

IHCC: PBZ\_KBN\_122/P05/2004.

IPOBCS: Liga Portuguesa Contra o Cancro.

kConFab: The National Breast Cancer Foundation, and previously by the National Health and Medical Research Council (NHMRC), the Queensland Cancer Fund, the Cancer Councils of New South Wales, Victoria, Tasmania and South Australia, and the Cancer Foundation of Western Australia. MAYO: NIH grants CA116167, CA192393 and CA176785, an NCI Specialized Program of Research Excellence (SPORE) in Breast Cancer (CA116201), and a grant from the Breast Cancer Research Foundation.

SWE-BCRA: the Swedish Cancer Society.

UPENN: Breast Cancer Research Foundation; Susan G. Komen Foundation for the cure, Basser Research Center for BRCA.

VFCTG: Victorian Cancer Agency, Cancer Australia, National Breast Cancer Foundation.

## Consortia

### kConFab Investigators

Amanda B. Spurdle<sup>1</sup>, Georgia Chenevix-Trench<sup>1</sup>, Jonathan Beesley<sup>1</sup>, Xiao Qing Chen<sup>1</sup>, Helene Holland<sup>1</sup>, Heather Thorne<sup>2</sup>, Stephen Fox<sup>2</sup>, Eveline NiederMayr<sup>2</sup>

### HEBON Investigators

Maartje J. Hoening<sup>3</sup>, Matti A. Rookus<sup>4</sup>, Frans B.L. Hogervorst<sup>5</sup>, Flora van Leeuwen<sup>5</sup>, Muriel A. Adank<sup>5</sup>, Denise Jenner<sup>5</sup>, Johanna M. Collée<sup>6</sup>, Ans M.W. van den Ouweland<sup>7</sup>, Ingrid A. Boere<sup>8</sup>, Christi J. van Asperen<sup>9</sup>, Peter Devilee<sup>10,11</sup>, Rob B. van der Luijt<sup>12</sup>, T.C.T.E.F. van Cronenburg<sup>13</sup>, M.R. Wevers<sup>14</sup>, Arjen R. Mensenkamp<sup>15</sup>, Margreet G.E.M. Ausems<sup>12</sup>, Marco Koudijs<sup>12</sup>, Theo A.M. van Os<sup>16</sup>, Klaartje van Engelen<sup>17</sup>,



Johan J.P. Gille<sup>17</sup>, Encarna B. Gómez García<sup>18</sup>, Marinus J. Blok<sup>19</sup>, Maaïke de Boer<sup>20</sup>, Lieke P.V. Berger<sup>21</sup>, Annemieke H. van der Hout<sup>22</sup>, Marian J.E. Mourits<sup>21</sup>, Geertruida H. de Bock<sup>21</sup>, Sabine Siesling<sup>23</sup>, Janneke Verloop<sup>23</sup>, Esther C. van den Broek<sup>24</sup>

#### **SWE-BRCA Investigators**

Ake Borg<sup>25</sup>, Håkan Olsson<sup>26</sup>, Helena Jernström<sup>27</sup>, Karin Henriksson<sup>28</sup>, Katja Harbst<sup>27</sup>, Maria Soller<sup>29</sup>, Ulf Kristoffersson<sup>27</sup>, Anna Öfverholm<sup>30</sup>, Margareta Nordling<sup>30</sup>, Per Karlsson<sup>31</sup>, Zakaria Einbeigi<sup>31</sup>, Anna von Wachenfeldt<sup>32</sup>, Annelie Liljegren<sup>32</sup>, Annika Lindblom<sup>33, 34</sup>, Brita Arver<sup>32</sup>, Gisela Barbany Bustinza<sup>34</sup>, Johanna Rantala<sup>35</sup>, Beatrice Melin<sup>36</sup>, Christina Edwinsdotter Ardnor<sup>37</sup>, Monica Emanuelsson<sup>36</sup>, Hans Ehrencrona<sup>29</sup>, Maritta Hellström-Pigg<sup>38</sup>, Richard Rosenquist<sup>38</sup>, Marie Stenmark-Askmal<sup>39</sup>, Sigrun Liedgren<sup>39</sup>

<sup>1</sup> QIMR Berghofer Medical Research Institute, Department of Genetics and Computational Biology, Brisbane, Queensland, Australia.

<sup>2</sup> Peter MacCallum Cancer Center, Research Department, Melbourne, Victoria, Australia.

<sup>3</sup> Erasmus MC Cancer Institute, Department of Medical Oncology, Family Cancer Clinic, Rotterdam, The Netherlands.

<sup>4</sup> The Netherlands Cancer Institute, Department of Epidemiology, Amsterdam, The Netherlands.

<sup>5</sup> The Netherlands Cancer Institute - Antoni van Leeuwenhoek hospital, Family Cancer Clinic, Amsterdam, The Netherlands.

<sup>6</sup> Erasmus University Medical Center, Rotterdam, The Netherlands.

<sup>7</sup> Erasmus University Medical Center, Department of Clinical Genetics, Rotterdam, The Netherlands.

<sup>8</sup> Erasmus MC Cancer Institute, Department of Medical Oncology, Rotterdam, The Netherlands.

<sup>9</sup> Leiden University Medical Center, Department of Clinical Genetics, Leiden, The Netherlands.

<sup>10</sup> Leiden University Medical Center, Department of Pathology, Leiden, The Netherlands.

<sup>11</sup> Leiden University Medical Center, Department of Human Genetics, Leiden, The Netherlands.

<sup>12</sup> University Medical Center, Department of Medical Genetics, Utrecht, The Netherlands.

<sup>13</sup> Leiden University Medical Center, Leiden, The Netherlands.

<sup>14</sup> Radboud University Medical Center, Nijmegen, The Netherlands.

<sup>15</sup> Radboud University Medical Center, Department of Human Genetics, Nijmegen, The Netherlands.

<sup>16</sup> Amsterdam UMC, location AMC, Department of Clinical Genetics, Amsterdam, The Netherlands.

<sup>17</sup> VU University Medical Center, Department of Clinical Genetics, Amsterdam, The Netherlands.

<sup>18</sup> Maastricht University Medical Center, Department of Clinical Genetics and GROW, School for Oncology and Developmental Biology, Maastricht, The Netherlands.

<sup>19</sup> Maastricht University Medical Center, Department of Clinical Genetics, Maastricht, The Netherlands.

<sup>20</sup> Maastricht University Medical Center, Maastricht, The Netherlands.

<sup>21</sup> University Medical Center Groningen, University Groningen, Groningen, The Netherlands.

<sup>22</sup> University Medical Center Groningen, University Groningen, Department of Genetics, Groningen, The Netherlands.

<sup>23</sup> Netherlands Comprehensive Cancer Organisation (IKNL), Utrecht, The Netherlands.

<sup>24</sup> Erasmus University, The nationwide network and registry of histo- and cytopathology in The Netherlands (PALGA), Rotterdam, The Netherlands.

<sup>25</sup> Lund University and Skåne University Hospital, Department of Oncology, Lund, Sweden.

<sup>26</sup> Lund University, Department of Cancer Epidemiology, Clinical Sciences, Lund, Sweden.

<sup>27</sup> Lund University, LUCC - Lund University Cancer Centre, Lund, Sweden.

<sup>28</sup> Lund University Hospital, Oncologic Centre, Regional Tumour Registry, Lund, Sweden.

<sup>29</sup> Lund University Hospital, Department of Clinical Genetics, Lund, Sweden.

<sup>30</sup> Sahlgrenska University Hospital, Department of Clinical Genetics, Gothenburg, Sweden.

<sup>31</sup> Sahlgrenska University Hospital, Department of Oncology, Gothenburg, Sweden.

<sup>32</sup> Karolinska Institutet, Department of Oncology, Stockholm, Sweden.

<sup>33</sup> Karolinska Institutet, Department of Molecular Medicine and Surgery, Stockholm, Sweden.

<sup>34</sup> Karolinska University Hospital, Department of Clinical Genetics, Stockholm, Sweden.

<sup>35</sup> Karolinska Institutet, Clinical Genetics, Stockholm, Sweden.

<sup>36</sup> Umeå University, Department of Radiation Sciences, Oncology, Umeå, Sweden.

<sup>37</sup> Changhua Christian Hospital, Department of Surgery, Changhua, Taiwan.

<sup>38</sup> Uppsala University, Department of Immunology, Genetics and Pathology, Uppsala, Sweden.

<sup>39</sup> Linköping University, Division of Clinical Genetics, Department of Clinical and Experimental Medicine, Faculty of Health Sciences, Linköping, Sweden.

**Supplementary Data 1.** Identification of potential target genes based on variant position and functional annotations. (separate .xlsx file)

**Supplementary Data 2.** eQTL-analysis in GTEx and Westra et al. data. (separate .xlsx file)